



# DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

## P03-07. Autologous Neutralizing Antibodies That Select Viral Escape Variants Emerge Late After SIV Infection of Rhesus Monkeys

The Harvard community has made this article openly available.  
[Please share](#) how this access benefits you. Your story matters.

Citation	Yeh, W. W., I. Rahman, P. Hraber, A. Giri, D. Nevidomskyte, R. T. Coffey, M. Asmal, et al. 2009. P03-07. Autologous neutralizing antibodies that select viral escape variants emerge late after SIV infection of rhesus monkeys. <i>Retrovirology</i> 6(Suppl 3): P24.
Published Version	<a href="https://doi.org/10.1186/1742-4690-6-S3-P24">doi:10.1186/1742-4690-6-S3-P24</a>
Accessed	February 19, 2015 8:12:07 AM EST
Citable Link	<a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:4875884">http://nrs.harvard.edu/urn-3:HUL.InstRepos:4875884</a>
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA">http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA</a>

*(Article begins on next page)*

Poster presentation

Open Access

## **P03-07. Autologous neutralizing antibodies that select viral escape variants emerge late after SIV infection of rhesus monkeys**

WW Yeh<sup>\*1</sup>, I Rahman<sup>1</sup>, P Hrabec<sup>2</sup>, A Giri<sup>1</sup>, D Nevidomskyte<sup>1</sup>, RT Coffey<sup>1</sup>, M Asmal<sup>1</sup>, S Miljkovic<sup>1</sup>, JB Whitney<sup>1</sup>, BF Keele<sup>3</sup>, GM Shaw<sup>3</sup>, BT Korber<sup>2</sup>, MS Seaman<sup>1</sup> and NL Letvin<sup>1</sup>

Address: <sup>1</sup>Division of Viral Pathogenesis, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, US, <sup>2</sup>Los Alamos National Laboratories, Los Alamos, NM, USA and <sup>3</sup>University of Alabama at Birmingham, Birmingham, AL, USA

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, **6**(Suppl 3):P24 doi:10.1186/1742-4690-6-S3-P24

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P24>

© 2009 Yeh et al; licensee BioMed Central Ltd.

### **Background**

Autologous neutralizing antibody responses against HIV-1 have been shown to emerge a number of months after primary infection. It is not known why neutralizing antibodies against HIV-1 are relatively slow to appear. We use the SIV/rhesus macaque model to characterize the kinetics of autologous neutralizing antibody responses and envelope sequence evolution in 4 rhesus monkeys infected intrarectally with SIVmac251.

### **Methods**

We assessed the presence of autologous neutralizing antibodies at 3, 5, 8, 16, and 22 months post-infection using pseudovirion-based, luciferase-reporter gene neutralizing antibody assays. We then analyzed full-length SIV env sequences at selected time points. The capacity of sera to neutralize variant viruses was also determined.

### **Results**

Sera from 4 SIVmac251-infected monkeys neutralized the inoculating virus at 8 months, but not at 5 months after infection. These sera did not neutralize autologous or heterologous SIV variants that were cloned at later time points, indicating that variant viruses had mutated to escape neutralization. After escape, these circulating viral variants became the predominant viral quasispecies in all animals. The majority of genotypic changes clustered in variable loops 1/2 (V1/2) and 4 (V4) of env. These

changes resulted in the deletion of neutralizing epitopes and shifts in potential N-linked glycosylation sites.

### **Conclusion**

We show that SIVmac251, a strain of SIV that is relatively resistant to antibody-mediated neutralization, induces potent autologous neutralizing antibodies by 8 months post-infection. The late emergence of neutralizing antibodies appears to be related to the inherent properties of primate lentiviruses, since the kinetics of SIVmac251 neutralization mirrors what has been reported with HIV-1. The evolution of SIV env implicates V1/2 and V4 region as the determinants of antibody neutralization. The development of these antibodies resulted in complete replacement of neutralization-sensitive virus by resistant viruses, providing evidence that humoral immune responses exert potent selection pressure on SIV despite their late emergence.